

© 2002 International Press
Adv. Theor. Math. Phys. **6** (2002) 1163–1182

Positive Mass Theorem on Manifolds admitting Corners along a Hypersurface

Pengzi Miao

Department of Mathematics
Stanford University
Stanford, CA 94305
USA
mpengzi@math.stanford.edu

Abstract

We study a class of non-smooth asymptotically flat manifolds on which metrics fail to be C^1 across a hypersurface Σ . We first give an approximation scheme to mollify the metric, then we show that the Positive Mass Theorem [8] still holds on these manifolds if a geometric boundary condition is satisfied by metrics separated by Σ .

1 Introduction and Statement of Results

The well-known Positive Mass Theorem in general relativity was first proved by R. Schoen and S.T. Yau in [8] for smooth asymptotically flat manifolds with non-negative scalar curvature. It is interesting to know on what kind of non-smooth Riemannian manifolds their techniques and results can be generalized. In this paper we study this question in a special setting where the metric fails to be C^1 across a hypersurface.

e-print archive: <http://lanl.arXiv.org/abs/math-ph/0212025>

Let $n \geq 3$ be a dimension for which the classical PMT [8] holds. Let $\alpha \in (0, 1)$ be a fixed number and M be an oriented n -dimensional smooth differentiable manifold with no boundary. We assume that there exists a compact domain $\Omega \subset M$ so that $M \setminus \Omega$ is diffeomorphic to $\mathbb{R}^n$ minus a ball and $\Sigma = \partial\Omega$ is a smooth hypersurface in M .

Definition 1. A metric $\mathcal{G}$ admitting corners along Σ is defined to be a pair of (g_-, g_+) , where g_- and g_+ are $C_{loc}^{2,\alpha}$ metrics on Ω and $M \setminus \overline{\Omega}$ so that they are C^2 up to the boundary and they induce the same metric on Σ .

Definition 2. Given $\mathcal{G} = (g_-, g_+)$, we say $\mathcal{G}$ is **asymptotically flat** if the manifold $(M \setminus \Omega, g_+)$ is asymptotically flat in the usual sense (see [7]).

Definition 3. The **mass** of $\mathcal{G} = (g_-, g_+)$ is defined to be the mass of g_+ (see [7]) whenever the latter exists.

One of our main motivation to study such a pair $\mathcal{G} = (g_-, g_+)$ is its implicit relation with Bartnik's quasi-local mass of the bounded Riemannian domain $(\overline{\Omega}, g_-)$. It is generally conjectured that there exists a g_+ on $M \setminus \Omega$ so that $\mathcal{G} = (g_-, g_+)$ is a minimal mass extension of $(\overline{\Omega}, g_-)$ in the sense of [1].

Under a geometric boundary condition which originated in [2], we prove the following Positive Mass Theorem for $\mathcal{G}$.

Theorem 1. Let $\mathcal{G} = (g_-, g_+)$ be an asymptotically flat metric admitting corners along Σ . Suppose that the scalar curvature of g_- , g_+ is non-negative in Ω , $M \setminus \overline{\Omega}$, and

$$H(\Sigma, g_-) \geq H(\Sigma, g_+), \quad (\text{H})$$

where $H(\Sigma, g_-)$ and $H(\Sigma, g_+)$ represent the mean curvature of Σ in $(\overline{\Omega}, g_-)$ and $(M \setminus \Omega, g_+)$ with respect to unit normal vectors pointing to the unbounded region. Then the mass of $\mathcal{G}$ is non-negative. Furthermore, if $H(\Sigma, g_-) > H(\Sigma, g_+)$ at some point on Σ , $\mathcal{G}$ has a strict positive mass.

Remark 1. Under our sign convention for the mean curvature, we have that $H(S^{n-1}, g_o) = n - 1$, where S^{n-1} is the unit sphere in $\mathbb{R}^n$ and g_o is the Euclidean metric.

One direct corollary of this theorem is that the boundary behavior of a metric g on $\overline{\Omega}$ imposes subtle restriction on the scalar curvature of g inside Ω . For instance, we have that

Corollary 1.1. There does not exist a metric g with non-negative scalar curvature on a standard ball $\overline{B} \subset \mathbb{R}^n$ so that ∂B is isometric to S^{n-1} and the mean curvature of ∂B in $(\overline{B}, g)$ is greater but not equal to $n - 1$.

Based on the work of H. Bray and F. Finster [4], we have a rigidity characterization of $\mathcal{G}$ when its mass is zero.

Theorem 2. *Let $n = 3$ and g_-, g_+ satisfy all the assumptions in Theorem 1. If g_- and g_+ are at least $C_{loc}^{3,\alpha}$, then the mass of g_+ being zero implies that g_- and g_+ are flat away from Σ and they induce the same second fundamental form on Σ . Hence, (Ω, g_-) and $(M \setminus \overline{\Omega}, g_+)$ together can be isometrically identified with the Euclidean space $(\mathbb{R}^3, g_o)$.*

To illustrate the relevance of Theorem 2 to the quasi-local mass of a bounded Riemannian domain, we mention the following corollary.

Corollary 1.2. *Let (M^3, g) be a manifold with non-negative scalar curvature, possibly with boundary. Let g_σ be a metric on S^2 so that there exist two isometric embeddings $\phi_1 : (S^2, g_\sigma) \rightarrow (\mathbb{R}^3, g_o)$ and $\phi_2 : (S^2, g_\sigma) \rightarrow (M^3, g)$, where $\phi_1(S^2), \phi_2(S^2)$ each bounds a compact region Ω_1, Ω_2 in $\mathbb{R}^3, M^3$ that has connected boundary. Then, if*

$$H(\phi_2(S^2), g) \geq H(\phi_1(S^2), g_o),$$

Ω_2 is isometric to Ω_1 . In particular, Ω_2 has trivial topology.

Remark 2. *If we replace S^2 by an arbitrary compact surface Σ_g with genus $g \geq 1$, under the same assumption, our argument still works to show that the region bounded by $\phi_2(\Sigma_g)$ is **flat**.*

2 Explanation of condition (H)

In this section we give a motivation for the geometric boundary condition (H). One will see that it can be interpreted as a statement that the scalar curvature of $\mathcal{G}$ is distributionally non-negative across Σ .

Let g be a C^2 metric in a tubular neighborhood of Σ and ν be a unit normal vector field to Σ . Let K be the Gaussian curvature of Σ with respect to the induced metric $g|_\Sigma$ and R be the scalar curvature of g . Taking trace of the Gauss equation, we have that

$$2K = R - 2Ric(\nu, \nu) + H^2 - |A|^2, \quad (1)$$

where $Ric(\nu, \nu)$ is the Ricci curvature of g along ν , H and A are the mean curvature and the second fundamental form of Σ .

Assuming that Σ evolves with speed ν , we have the following evolution formula of the mean curvature

$$D_\nu H = -Ric(\nu, \nu) - |A|^2. \quad (2)$$

It follows from (1) and (2) that

$$R = 2K - (|A|^2 + H^2) - 2D_\nu H, \quad (3)$$

which suggests that $D_\nu H$ plays a dominant role in determining the sign of R if K, H and A are known to be bounded. In particular, for a metric $\mathcal{G} = (g_-, g_+)$ with $H(\Sigma, g_-) > H(\Sigma, g_+)$, the scalar curvature of $\mathcal{G}$ across Σ looks like a positive Dirac-Delta function with support in Σ . Hence, the spirit of Theorem 1 is that PMT still holds even if the scalar curvature is only assumed to be distributionally non-negative across Σ .

Remark 3. *The geometric boundary condition (H) was first introduced by R. Bartnik in [2], where he suggested the static metric extension conjecture for a bounded domain in a time-symmetric initial data set.*

3 Smoothing $\mathcal{G}$ across Σ

Given $\mathcal{G} = (g_-, g_+)$ on M , we want to approximate $\mathcal{G}$ by metrics which are C^2 across Σ .

First, we use the Gaussian coordinates of Σ to modify the differential structure on M so that $\mathcal{G}$ becomes a continuous metric across Σ . Let $U_-^{2\epsilon}$ be a 2ϵ -tubular neighborhood of Σ in $(\overline{\Omega}, g_-)$ for some $\epsilon > 0$. Let

$$\Phi_- : \Sigma \times (-2\epsilon, 0] \longrightarrow U_-^{2\epsilon}$$

be a diffeomorphism so that the pull back metric $\Phi_-^*(g_-)$ has the form $\Phi_-^*(g_-) = g_{-ij}(x, t)dx^i dx^j + dt^2$, where t is the coordinate for $(-2\epsilon, 0]$, $(x^1, \dots, x^{n-1})$ are local coordinates for Σ and i, j runs through $1, \dots, n-1$. Similarly, we have $\Phi_+ : \Sigma \times [0, 2\epsilon) \longrightarrow U_+^{2\epsilon}$, where $U_+^{2\epsilon}$ is a 2ϵ -tubular neighborhood of Σ in $(M \setminus \Omega, g_+)$ and $\Phi_+^*(g_+) = g_{+ij}(x, t)dx^i dx^j + dt^2$. Identifying $U = U_-^{2\epsilon} \cup U_+^{2\epsilon}$ with $\Sigma \times (-2\epsilon, 2\epsilon)$, we define $\tilde{M}$ to be a possibly new differentiable manifold with the background topological space M and the differential structure determined by the open covering $\{\Omega, M \setminus \overline{\Omega}, U\}$. Since $g_-|_\Sigma = g_+|_\Sigma$, $\mathcal{G}$ becomes a continuous metric g on $\tilde{M}$. Inside $U = \Sigma \times (-2\epsilon, 2\epsilon)$, we have that

$$g = g_{ij}(x, t)dx^i dx^j + dt^2, \quad (4)$$

where $g_{ij}(x, t) = g_{-ij}(x, t)$ when $t \leq 0$ and $g_{ij}(x, t) = g_{+ij}(x, t)$ when $t \geq 0$.

Second, we mollify the metric g inside U . Let $i \in \{0, 1, 2\}$, we define $\mathcal{S}^i(\Sigma)$ to be the Banach space of C^i symmetric $(0, 2)$ tensors on Σ equipped with the usual C^i norm and $\mathcal{M}^i(\Sigma)$ to be the open and convex subset of

$\mathcal{S}^i(\Sigma)$ consisting of C^i metrics. By (4) we have a well defined path in each $\mathcal{M}^i(\Sigma)$,

$$\gamma : (-2\epsilon, 2\epsilon) \longrightarrow \mathcal{M}^2(\Sigma) \hookrightarrow \mathcal{M}^1(\Sigma) \hookrightarrow \mathcal{M}^0(\Sigma) \quad (5)$$

where

$$\gamma(t) = g_{ij}(x, t) dx^i dx^j. \quad (6)$$

By assumption, γ is a continuous path in $\mathcal{M}^2(\Sigma)$ and a piecewise C^1 path in $\mathcal{M}^1(\Sigma)$. Hence, there exists $L > 0$ depending only on $\mathcal{G}$ such that

$$\|\gamma(t) - \gamma(s)\|_{\mathcal{M}^1(\Sigma)} \leq L|t - s|, \quad \forall s, t \in [-\frac{3}{2}\epsilon, \frac{3}{2}\epsilon]. \quad (7)$$

We choose $\phi(t) \in C_c^\infty([-1, 1])$ to be a standard mollifier on $\mathbb{R}^1$ such that

$$0 \leq \phi \leq 1 \quad \text{and} \quad \int_{-1}^1 \phi(t) dt = 1. \quad (8)$$

Let $\sigma(t) \in C_c^\infty([-\frac{1}{2}, \frac{1}{2}])$ be another cut-off function such that

$$\begin{cases} 0 \leq \sigma(t) \leq \frac{1}{100} & t \in \mathbb{R}^1 \\ \sigma(t) = \frac{1}{100} & |t| < \frac{1}{4} \\ 0 < \sigma(t) \leq \frac{1}{100} & \frac{1}{4} < |t| < \frac{1}{2}. \end{cases} \quad (9)$$

Given any $0 < \delta \ll \epsilon$, let

$$\sigma_\delta(t) = \delta^2 \sigma\left(\frac{t}{\delta}\right) \quad (10)$$

and we define

$$\begin{aligned} \gamma_\delta(s) &= \int_{\mathbb{R}^1} \gamma(s - \sigma_\delta(s)t) \phi(t) dt, & s \in (-\epsilon, \epsilon) \\ &= \begin{cases} \int_{\mathbb{R}^1} \gamma(t) \left(\frac{1}{\sigma_\delta(s)} \phi\left(\frac{s-t}{\sigma_\delta(s)}\right) \right) dt, & \sigma_\delta(s) > 0 \\ \gamma(s), & \sigma_\delta(s) = 0, \end{cases} \end{aligned} \quad (11)$$

where the integral takes place in $\mathcal{S}^0(\Sigma)$. By the convexity of $\mathcal{M}^0(\Sigma)$ in $\mathcal{S}^0(\Sigma)$, γ_δ is a path in $\mathcal{M}^0(\Sigma)$. We have the following elementary lemmas concerning the property of γ_δ and its relation with γ .

Lemma 3.1. $\gamma_\delta(s)$ is a C^2 path in $\mathcal{M}^0(\Sigma)$ and a C^1 path in $\mathcal{M}^1(\Sigma)$.

Proof: Let $i \in \{1, 2\}$. The fact that $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^{2-i}(\Sigma)$ is C^i away from 0 implies that $\gamma_\delta : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^{2-i}(\Sigma)$ is C^i away from $[-\frac{\delta^2}{100}, \frac{\delta^2}{100}]$. For $s \in (-\frac{\delta}{4}, \frac{\delta}{4})$, $\sigma_\delta(s) = \frac{\delta^2}{100}$ and (11) turns into

$$\gamma_\delta(s) = \int_{\mathbb{R}^1} \gamma(t) \left(\frac{1}{(\frac{\delta^2}{100})} \phi\left(\frac{s-t}{(\frac{\delta^2}{100})}\right) \right) dt, \quad (12)$$

which is the standard mollification of γ by ϕ with a constant scaling factor $\frac{\delta^2}{100}$. Hence, $\gamma_\delta(s) : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^{2-i}(\Sigma)$ is smooth in $(-\frac{\delta}{4}, \frac{\delta}{4})$. $\square$

Lemma 3.2. $\gamma_\delta(s)$ is a C^0 path in $\mathcal{M}^2(\Sigma)$ which is uniformly close to γ and agrees with γ outside $(-\frac{\delta}{2}, \frac{\delta}{2})$.

Proof: The continuity of $\gamma_\delta : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^2(\Sigma)$ follows directly from that of γ . The estimate

$$\begin{aligned} \|\gamma_\delta(s) - \gamma(s)\|_{\mathcal{M}^2(\Sigma)} &= \left\| \int_{\mathbb{R}^1} (\gamma(s - \sigma_\delta(s)t) - \gamma(s)) \phi(t) dt \right\|_{\mathcal{M}^2(\Sigma)} \\ &\leq \int_{\mathbb{R}^1} \|\gamma(s - \sigma_\delta(s)t) - \gamma(s)\|_{\mathcal{M}^2(\Sigma)} \phi(t) dt \end{aligned} \quad (13)$$

shows it is uniformly close to γ . Finally, $\sigma_\delta(s) = 0$ for $|s| > \frac{\delta}{2}$ implies

$$\gamma_\delta(s) = \int_{\mathbb{R}^1} \gamma(s) \phi(t) dt = \gamma(s). \quad (14)$$

□

Lemma 3.3. $\|\gamma_\delta(s) - \gamma(s)\|_{\mathcal{M}^0(\Sigma)} \leq L\delta^2$, for $s \in (-\epsilon, \epsilon)$.

Proof: It follows from (7) that

$$\begin{aligned} \|\gamma_\delta(s) - \gamma(s)\|_{\mathcal{M}^0(\Sigma)} &\leq \int_{\mathbb{R}^1} \|\gamma(s - \sigma_\delta(s)t) - \gamma(s)\|_{\mathcal{M}^0(\Sigma)} \phi(t) dt \\ &\leq L\delta^2. \end{aligned} \quad (15)$$

□

Now we define

$$g_\delta = \begin{cases} \gamma_\delta(t) + dt^2, & (x, t) \in \Sigma \times (-\epsilon, \epsilon) \\ g, & (x, t) \notin \Sigma \times (-\epsilon, \epsilon). \end{cases} \quad (16)$$

Lemma 3.1, 3.2 and 3.3 imply that g_δ is a globally C^2 metric on $\tilde{M}$ which agrees with g outside a strip region $\Sigma \times (-\frac{\delta}{2}, \frac{\delta}{2})$ and is uniformly close to g in C^0 topology.

Next, we proceed to estimate the scalar curvature of g_δ . We use the notations defined in section 2 with a lower index δ to denote the corresponding quantities of g_δ . By (16), the vector field $\frac{\partial}{\partial t}$ is perpendicular to the slice $\Sigma \times \{t\}$ for each $t \in (-\epsilon, \epsilon)$. Hence, inside $\Sigma \times (-\epsilon, \epsilon)$, (3) shows that

$$R_\delta(x, t) = 2K_\delta(x, t) - (|A_\delta(x, t)|^2 + H_\delta(x, t)^2) - 2\frac{\partial}{\partial t}H_\delta(x, t). \quad (17)$$

We will estimate each term on the right of (17). First we note that $K_\delta(x, t)$ is determined only by $\gamma_\delta(t)$, Lemma 3.2 then implies that $K_\delta(x, t)$ is bounded by constants depending only on $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^2(\Sigma)$.

To estimate $A_\delta(x, t)$ and $H_\delta(x, t)$, we compute the first order derivative of $\gamma_\delta : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$ because of the definition

$$A_{\delta ij}(x, t) = \langle \frac{\partial}{\partial x^j}, \nabla_{\frac{\partial}{\partial x^i}}^\delta \partial_t \rangle = \frac{1}{2} \frac{\partial}{\partial t} \gamma_{\delta ij}(x, t). \quad (18)$$

By (11) we have that

$$\frac{\partial}{\partial t} \gamma_{\delta ij}(x, t) = \frac{\partial}{\partial t} \int_{\mathbb{R}^1} \gamma_{ij}(t - \sigma_\delta(t)s) \phi(s) ds. \quad (19)$$

When $|t| > \frac{\delta^2}{100}$, (19) gives that

$$\begin{aligned} \frac{\partial}{\partial t} \gamma_{\delta ij}(x, t) &= \int_{\mathbb{R}^1} \frac{d}{dt} \{ \gamma_{ij}(t - \sigma_\delta(t)s) \} \phi(s) ds \\ &= \int_{\mathbb{R}^1} \gamma'_{ij}(t - \sigma_\delta(t)s) \{ 1 - s \delta \sigma'(\frac{t}{\delta}) \} \phi(s) ds. \end{aligned} \quad (20)$$

When $|t| < \frac{\delta}{4}$, (12) implies that

$$\begin{aligned} \frac{\partial}{\partial t} \gamma_{\delta ij}(x, t) &= \frac{\partial}{\partial t} \int_{\mathbb{R}^1} \gamma_{ij}(s) \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds \\ &= \int_{\mathbb{R}^1} \gamma_{ij}(s) \frac{d}{dt} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds \\ &= (-1) \int_{\mathbb{R}^1} \gamma_{ij}(s) \frac{d}{ds} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds \\ &= (-1) \int_{-\infty}^0 \gamma_{ij}(s) \frac{d}{ds} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds \\ &\quad + (-1) \int_0^\infty \gamma_{ij}(s) \frac{d}{ds} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds. \end{aligned} \quad (21)$$

Integrating by parts and considering the fact $\gamma(t)$ is continuous at 0, we have that

$$\begin{aligned} \frac{\partial}{\partial t} \gamma_{\delta ij}(x, t) &= \int_{\mathbb{R}^1} \gamma'_{ij}(s) \left\{ \frac{100}{\delta^2} \phi\left(\frac{100(t-s)}{\delta^2}\right) \right\} ds \\ &= \int_{\mathbb{R}^1} \gamma'_{ij}(t - \sigma_\delta(t)s) \phi(s) ds. \end{aligned} \quad (22)$$

Therefore, for every $t \in (-\epsilon, \epsilon)$, we have that

$$\frac{\partial}{\partial t} \gamma_{\delta ij}(x, t) = \int_{\mathbb{R}^1} \gamma'_{ij}(t - \sigma_\delta(t)s) \left\{ 1 - s \delta \sigma'(\frac{t}{\delta}) \right\} \phi(s) ds, \quad (23)$$

which shows that $A_\delta(x, t)$ is bounded by constants depending only on $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$. Since $H_\delta(x, t) = g_\delta^{ij} A_{\delta ij}$, Lemma 3.2 and (23) also imply

that $H_\delta(x, t)$ is bounded by constants depending only on $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$.

To estimate $\frac{\partial}{\partial t} H_\delta(x, t)$, we need to compute the second order derivative of $\gamma_\delta : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$. A similar calculation as above gives that, for $|t| > \frac{\delta^2}{100}$,

$$\begin{aligned} \frac{\partial^2}{\partial^2 t} \gamma_{\delta ij}(x, t) &= \int_{\mathbb{R}^1} \gamma''_{ij}(t - \sigma_\delta(t)s) \left\{ 1 - s\delta\sigma'(\frac{t}{\delta}) \right\}^2 \phi(s) ds + \\ &\quad \int_{\mathbb{R}^1} \gamma'_{ij}(t - \sigma_\delta(t)s) \left\{ -s\sigma''(\frac{t}{\delta}) \right\} \phi(s) ds \end{aligned} \quad (24)$$

and, for $|t| < \frac{\delta}{4}$,

$$\begin{aligned} \frac{\partial^2}{\partial^2 t} \gamma_{\delta ij}(x, t) &= \int_{\mathbb{R}^1} \gamma''_{ij}(t - \sigma_\delta(t)s) \phi(s) ds + \\ &\quad \{g_{+ij}'(0) - g_{-ij}'(0)\} \left\{ \frac{100}{\delta^2} \phi(\frac{100t}{\delta^2}) \right\}. \end{aligned} \quad (25)$$

Since

$$\frac{\partial}{\partial t} H_\delta(x, t) = \frac{\partial}{\partial t} \{g_\delta^{ij}(x, t)\} A_{\delta ij}(x, t) + g_\delta^{ij}(x, t) \frac{\partial}{\partial t} A_{\delta ij}(x, t), \quad (26)$$

(24), (23) and Lemma 3.2 imply that, outside $\Sigma \times [-\frac{100}{\delta^2}, \frac{100}{\delta^2}]$, $\frac{\partial}{\partial t} H_\delta(x, t)$ is bounded by constants only depending on $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$. On the other hand, inside $\Sigma \times [-\frac{\delta^2}{100}, \frac{\delta^2}{100}]$, (25) and (26) show that

$$\begin{aligned} \frac{\partial}{\partial t} H_\delta(x, t) &= \frac{\partial}{\partial t} \{g_\delta^{ij}(x, t)\} A_{\delta ij}(x, t) + \\ &\quad \frac{1}{2} g_\delta^{ij}(x, t) \left\{ \int_{\mathbb{R}^1} \gamma''_{ij}(t - \sigma_\delta(t)s) \phi(s) ds \right\} + \\ &\quad \frac{1}{2} g_\delta^{ij}(x, t) \{g_{+ij}'(0) - g_{-ij}'(0)\} \left\{ \frac{100}{\delta^2} \phi(\frac{100t}{\delta^2}) \right\}. \end{aligned} \quad (27)$$

The first two terms on the right are bounded by constants depending only on $\gamma : (-\epsilon, \epsilon) \longrightarrow \mathcal{M}^0(\Sigma)$. For the third one, we rewrite it as

$$\begin{aligned} &\frac{1}{2} \{g_\delta^{ij}(x, t) - g^{ij}(x, 0)\} \{g_{+ij}'(0) - g_{-ij}'(0)\} \left\{ \frac{100}{\delta^2} \phi(\frac{100t}{\delta^2}) \right\} \\ &\quad + \{H(\Sigma, g_+)(x) - H(\Sigma, g_-)(x)\} \left\{ \frac{100}{\delta^2} \phi(\frac{100t}{\delta^2}) \right\}. \end{aligned} \quad (28)$$

By (7), Lemma 3.3 and the fact $|t| \leq \frac{\delta^2}{100}$, we have that

$$\begin{aligned} |g_\delta^{ij}(x, t) - g^{ij}(x, 0)| &\leq |g_\delta^{ij}(x, t) - g^{ij}(x, t)| + |g^{ij}(x, t) - g^{ij}(x, 0)| \\ &\leq CL\delta^2 + CL\delta^2, \end{aligned} \quad (29)$$

where $C > 0$ only depends on $\mathcal{G}$. Hence, we conclude that

$$\frac{\partial}{\partial t} H_\delta(x, t) = O(1) + \{H(\Sigma, g_+)(x) - H(\Sigma, g_-)(x)\} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100t}{\delta^2}\right) \right\} \quad (30)$$

inside $\Sigma \times [-\frac{\delta^2}{100}, \frac{\delta^2}{100}]$, where $O(1)$ represents bounded quantities with bounds depending only on $\mathcal{G}$.

We summarize the features of $\{g_\delta\}$ in the following proposition.

Proposition 3.1. *Let $\mathcal{G} = (g_-, g_+)$ be a metric admitting corners along Σ . Then $\exists$ a family of C^2 metrics $\{g_\delta\}_{0 < \delta \leq \delta_0}$ on $\tilde{M}$ so that g_δ is uniformly close to g on $\tilde{M}$, $g_\delta = g$ outside $\Sigma \times (-\frac{\delta}{2}, \frac{\delta}{2})$ and the scalar curvature of g_δ satisfies*

$$R_\delta(x, t) = O(1), \quad \text{for } (x, t) \in \Sigma \times \left\{ \frac{\delta^2}{100} < |t| \leq \frac{\delta}{2} \right\} \quad (31)$$

$$R_\delta(x, t) = O(1) + \{H(\Sigma, g_-)(x) - H(\Sigma, g_+)(x)\} \left\{ \frac{100}{\delta^2} \phi\left(\frac{100t}{\delta^2}\right) \right\},$$

$$\text{for } (x, t) \in \Sigma \times \left[-\frac{\delta^2}{100}, \frac{\delta^2}{100}\right], \quad (32)$$

where $O(1)$ represents quantities that are bounded by constants depending only on $\mathcal{G}$, but not on δ .

In case $H(\Sigma, g_-) \equiv H(\Sigma, g_+)$, the following corollary generalizes a reflecting argument used by H. Bray in his proof of the Riemannian Penrose Inequality [3].

Corollary 3.1. *Given $\mathcal{G} = (g_-, g_+)$, if $H(\Sigma, g_-) \equiv H(\Sigma, g_+)$, then $\exists$ a family of C^2 metrics $\{g_\delta\}_{0 < \delta \leq \delta_0}$ on $\tilde{M}$ so that g_δ is uniformly close to g on $\tilde{M}$, $g_\delta = g$ outside $\Sigma \times (-\frac{\delta}{2}, \frac{\delta}{2})$ and the scalar curvature of g_δ is uniformly bounded inside $\Sigma \times [-\frac{\delta}{2}, \frac{\delta}{2}]$ with bounds depending only on $\mathcal{G}$, but not on δ .*

4 Proof of Theorem 1

We fix the following notations. Given a function f , we let f_+ and f_- denote the positive and negative part of f , i.e. $f = f_+ - f_-$ and $|f| = f_+ + f_-$. Given a metric g , we define the conformal Laplacian of g to be $L_g(u) = \Delta_g u - c_n R(g)u$, where $c_n = \frac{n-2}{4(n-1)}$ and $R(g)$ is the scalar curvature of g . The mass of g will be denoted by $m(g)$ if it exists. Finally, we let $C_0, C_1, C_2, \dots$ represent constants depending only on $\mathcal{G}$.

Throughout this section, we assume that $R(g_-), R(g_+) \geq 0$ in Ω , $M \setminus \overline{\Omega}$, and $H(\Sigma, g_-)(x) \geq H(\Sigma, g_+)(x)$ for all $x \in \Sigma$.

4.1 Conformal Deformation

We want to modify $\{g_\delta\}$ on $\tilde{M}$ to get C^2 metrics with non-negative scalar curvature. For that purpose we use conformal deformation. The following fundamental lemma is due to Schoen and Yau. Interested readers may refer to [8] for a detailed proof.

Lemma 4.1. [8] *Let g be a C^2 asymptotically flat metric on $\tilde{M}$ and f be a function that has the same decay rate at ∞ as $R(g)$, then $\exists$ a number $\epsilon_0 > 0$ depending only on the C^0 norm of g and the decay rate of g , ∂g and $\partial\partial g$ at ∞ so that if*

$$\left\{ \int_{\tilde{M}} |f_-|^{\frac{n}{2}} dg \right\}^{\frac{2}{n}} < \epsilon_0, \quad (33)$$

then

$$\begin{cases} \Delta_g u - c_n f u &= 0 \\ \lim_{x \rightarrow \infty} u &= 1 \end{cases} \quad (34)$$

has a C^2 positive solution u defined on $\tilde{M}$ so that $u = 1 + \frac{A}{|x|^{n-2}} + \omega$ for some constant A and some function ω , where $\omega = O(|x|^{1-n})$ and $\partial\omega = O(|x|^{-n})$.

For each δ , we consider the following equation

$$\begin{cases} \Delta_{g_\delta} u_\delta + c_n R_{\delta-} u_\delta &= 0 \\ \lim_{x \rightarrow \infty} u_\delta &= 1. \end{cases} \quad (35)$$

It follows from Proposition 3.1 and assumptions on $R(g_-)$ and $R(g_+)$ that

$$\begin{cases} R_{\delta-} = 0, & \text{outside } \Sigma \times [-\frac{\delta}{2}, \frac{\delta}{2}] \\ |R_{\delta-}| \leq C_0, & \text{inside } \Sigma \times [-\frac{\delta}{2}, \frac{\delta}{2}]. \end{cases} \quad (36)$$

Therefore, (33) holds with f and g replaced by $-R_{\delta-}$ and g_δ , for sufficiently small δ . We note that ϵ_0 can be chosen to be independent on δ because of Proposition 3.1. Hence the solution to (35) exists by Lemma 4.1. We have the following L^∞ and $C^{2,\alpha}$ estimate for $\{u_\delta\}$.

Proposition 4.1. $\lim_{\delta \rightarrow 0} \|u_\delta - 1\|_{L^\infty(\tilde{M})} = 0$ and $\|u_\delta\|_{C^{2,\alpha}(K)} \leq C_K$. Here K is any compact set in $\tilde{M} \setminus \Sigma$ and C_K only depends on g and K .

Proof: It suffices to obtain the L^∞ estimate of $|u_\delta - 1|$ because, once it is established, the $C^{2,\alpha}$ estimate will follow directly from the fact $\Delta_g u_\delta = 0$ outside $\Sigma \times [-\frac{\delta}{2}, \frac{\delta}{2}]$ and the standard Schauder theory. Let $w_\delta = u_\delta - 1$, we have that

$$\Delta_{g_\delta} w_\delta + c_n R_{\delta-} w_\delta = -c_n R_{\delta-} \quad (37)$$

where $w_\delta = \frac{A_\delta}{|x|^{n-2}} + \omega_\delta$ for some constant A_δ and some function ω_δ with the decay rate in Lemma 4.1. Multiply (37) by w_δ and integrate over $\tilde{M}$,

$$\int_{\tilde{M}} (w_\delta \triangle_{g_\delta} w_\delta + c_n R_{\delta-} w_\delta^2) dg_\delta = \int_{\tilde{M}} -c_n R_{\delta-} w_\delta dg_\delta. \quad (38)$$

Integrating by parts and using Hölder Inequality, we have that

$$\begin{aligned} \int_{\tilde{M}} |\nabla_{g_\delta} w_\delta|^2 dg_\delta &\leq c_n \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{n}{2}} dg_\delta \right)^{\frac{2}{n}} \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} \\ &\quad + c_n \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{2n}{n+2}} dg_\delta \right)^{\frac{n+2}{2n}} \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{2n}} \end{aligned} \quad (39)$$

On the other hand, the Sobolev Inequality gives that

$$\left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} \leq C_\delta \int_{\tilde{M}} |\nabla_{g_\delta} w_\delta|^2 dg_\delta, \quad (40)$$

where C_δ denotes the Sobolev Constant of the metric g_δ . It follows from (39), (40) and the elementary inequality $ab \leq \frac{a^2}{2} + \frac{b^2}{2}$ that

$$\begin{aligned} \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} &\leq C_\delta c_n \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{n}{2}} dg_\delta \right)^{\frac{2}{n}} \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} \\ &\quad + \frac{1}{2} C_\delta^2 c_n^2 \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{2n}{n+2}} dg_\delta \right)^{\frac{n+2}{n}} \\ &\quad + \frac{1}{2} \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}}. \end{aligned} \quad (41)$$

We note that Proposition 3.1 implies that C_δ is uniformly close to the Sobolev Constant of g . Hence, for sufficiently small δ , (41) gives that

$$\left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} \leq C \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{2n}{n+2}} dg_\delta \right)^{\frac{n+2}{n}} = o(1), \quad \text{as } \delta \rightarrow 0. \quad (42)$$

This $L^{\frac{2n}{n-2}}$ estimate and (37) imply the supremum estimate for w_δ

$$\begin{aligned} \sup_{\tilde{M}} |w_\delta| &\leq C \left\{ \left(\int_{\tilde{M}} w_\delta^{\frac{2n}{n-2}} dg_\delta \right)^{\frac{n-2}{2n}} + \left(\int_{\tilde{M}} |R_{\delta-}|^{\frac{n}{n-2}} dg_\delta \right)^{\frac{n-2}{n}} \right\} \\ &= o(1) \quad \text{as } \delta \rightarrow 0 \end{aligned} \quad (43)$$

by the standard linear theory (Theorem 8.17 in [5]). $\square$

Now we define

$$\tilde{g}_\delta = u_\delta^{\frac{4}{n-2}} g_\delta. \quad (44)$$

It follows from Proposition 4.1 that, passing to a subsequence, $\{\tilde{g}_\delta\}$ converges to g in C^0 topology on $\tilde{M}$ and in C^2 topology on compact sets away from Σ . By the conformal transformation formulae of scalar curvature [7], we also have that

$$\tilde{R}_\delta = -c_n^{-1} u_\delta^{-(\frac{n+2}{n-2})} L_{g_\delta}(u_\delta) = u_\delta^{\frac{4}{2-n}} R_{\delta+} \geq 0, \quad (45)$$

where $\tilde{R}_\delta$ represents the scalar curvature of $\tilde{g}_\delta$.

Lemma 4.2. *The mass of $\tilde{g}_\delta$ converges to the mass of $\mathcal{G}$.*

Proof: A straightforward calculation using the definition of mass reveals that

$$m(\tilde{g}_\delta) = m(g_\delta) + (n-1)A_\delta, \quad (46)$$

where A_δ is given by the expansion $u_\delta(x) = 1 + A_\delta|x|^{2-n} + O(|x|^{1-n})$. Applying integration by parts to (35) multiplied by u_δ , we have that

$$(2-n)\omega_n A_\delta = \int_{\tilde{M}} [|\nabla_{g_\delta} u_\delta|^2 - c_n R_{\delta-} u_\delta^2] dg_\delta, \quad (47)$$

where ω_n is the volume of the $n-1$ dimensional unit sphere in $\mathbb{R}^n$. It follows from that (46) and (47) that

$$m(g_\delta) = m(\tilde{g}_\delta) + \frac{n-1}{n-2}\omega_n \int_{\tilde{M}} [|\nabla_{g_\delta} u_\delta|^2 - c_n R_{\delta-} u_\delta^2] dg_\delta. \quad (48)$$

We note that the integral term above approaches 0 by Proposition 4.1, (36) and (39). Hence, we have that

$$\lim_{\delta \rightarrow 0} m(\tilde{g}_\delta) = \lim_{\delta \rightarrow 0} m(g_\delta) = m(\mathcal{G}).$$

□

Applying the classical PMT [8] to each $\tilde{g}_\delta$, we have that $m(\tilde{g}_\delta) \geq 0$. Thus, the non-negativity of $m(\mathcal{G})$ follows directly from Lemma 4.2.

4.2 Scalar Curvature Concentration

In this subsection, we assume that there exists strict jump of mean curvature across Σ , i.e.

$$H(\Sigma, g_-)(x) > H(\Sigma, g_+)(x) \quad \text{for some } x \in \Sigma.$$

We will prove that $\mathcal{G}$ has a strict positive mass.

Since $H(\Sigma, g_-)$ and $H(\Sigma, g_+)$ both are continuous functions on Σ , we can choose a compact set $K \subset \Sigma$ so that

$$H(\Sigma, g_-)(x) - H(\Sigma, g_+)(x) \geq \eta, \quad \forall x \in K \quad (49)$$

for some fixed $\eta > 0$. By Proposition 3.1, we have that

$$R_{\delta+}(x, t) \geq \eta \left\{ \frac{100}{\delta^2} \phi\left(\frac{100t}{\delta^2}\right) \right\} - C_0, \quad \forall (x, t) \in K \times \left[-\frac{\delta^2}{100}, \frac{\delta^2}{100}\right], \quad (50)$$

which suggests that the scalar curvature of g_δ and $\tilde{g}_\delta$ has a fixed amount of concentration on K .

To exploit this fact, we use conformal deformation again to make $\tilde{g}_\delta$ even scalar flat. Since $\tilde{R}_\delta = u_\delta^{\frac{4}{2-n}} R_{\delta+} \geq 0$, $\exists$ a C^2 positive solution to the following equations

$$\begin{cases} \Delta_{\tilde{g}_\delta} v_\delta - c_n \tilde{R}_\delta v_\delta &= 0 \\ \lim_{x \rightarrow \infty} v_\delta &= 1. \end{cases} \quad (51)$$

By the maximum principle, we have that

$$0 < v_\delta \leq 1. \quad (52)$$

Now define

$$\hat{g}_\delta = v_\delta^{\frac{4}{n-2}} \tilde{g}_\delta. \quad (53)$$

Similar to previous discussion, we know that $\hat{g}_\delta$ is an asymptotically flat metric and the scalar curvature of $\hat{g}_\delta$ is identically zero. Furthermore, $m(\hat{g}_\delta)$ and $m(\tilde{g}_\delta)$ are related by

$$m(\tilde{g}_\delta) = m(\hat{g}_\delta) + \frac{n-1}{n-2} \omega_n \int_{\tilde{M}} \left[|\nabla_{\tilde{g}_\delta} v_\delta|^2 + c_n \tilde{R}_\delta v_\delta^2 \right] d\tilde{g}_\delta, \quad (54)$$

where $m(\hat{g}_\delta) \geq 0$ by the classical PMT. Hence, to prove $m(\mathcal{G}) > 0$, it suffices to show the integral term in (54) has a strict positive lower bound.

Proposition 4.2.

$$\inf_{\delta > 0} \left\{ \int_{\tilde{M}} \left[|\nabla_{\tilde{g}_\delta} v_\delta|^2 + c_n \tilde{R}_\delta v_\delta^2 \right] d\tilde{g}_\delta \right\} > 0 \quad (55)$$

Proof: Assume (55) is not true, passing to a subsequence, we may assume that

$$\lim_{\delta \rightarrow 0} \int_{\tilde{M}} \left[|\nabla_{\tilde{g}_\delta} v_\delta|^2 + c_n \tilde{R}_\delta v_\delta^2 \right] d\tilde{g}_\delta = 0. \quad (56)$$

Since $\tilde{R}_\delta \geq 0$, (56) is equivalent to

$$\lim_{\delta \rightarrow 0} \int_{\tilde{M}} |\nabla_{\tilde{g}_\delta} v_\delta|^2 d\tilde{g}_\delta = 0 \quad \text{and} \quad \lim_{\delta \rightarrow 0} \int_{\tilde{M}} \tilde{R}_\delta v_\delta^2 d\tilde{g}_\delta = 0. \quad (57)$$

Outside $\Sigma \times [-\frac{\delta}{2}, \frac{\delta}{2}]$, we have $g_\delta = g$. Hence, (51) becomes

$$\Delta_{\tilde{g}_\delta} v_\delta - c_n \left(u_\delta^{\frac{4}{2-n}} R(g)_+ \right) v_\delta = 0. \quad (58)$$

It follows from Proposition 4.1, (52) and Schauder Estimates that, passing to a subsequence, v_δ converges to a function v in C^2 topology on compact sets away from Σ . By (57), we have that

$$\int_{\tilde{M} \setminus \Sigma} |\nabla_g v|^2 dg = 0, \quad (59)$$

which shows that v is a constant on Ω and $\tilde{M} \setminus \bar{\Omega}$.

We claim that $v = 1$ on $\tilde{M} \setminus \bar{\Omega}$. If not, we may assume $v = \beta < 1$ by (52). We fix a $\delta_0 \in (0, \epsilon)$ and denote the region inside $\Sigma \times \{\delta_0\}$ by Ω_{δ_0} . For each $\delta < \delta_0$, we let w_δ be the solution to the following equations

$$\begin{cases} \Delta_{\tilde{g}_\delta} w_\delta = 0 & \text{on } \tilde{M} \setminus \bar{\Omega}_{\delta_0} \\ w_\delta = v_\delta & \text{on } \Sigma \times \{\delta_0\} \\ w_\delta(x) \rightarrow 1 & \text{at } \infty. \end{cases} \quad (60)$$

Since w_δ minimizes the Dirichlet energy among all functions with the same boundary values, we have that

$$\int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_{\tilde{g}_\delta} w_\delta|^2 d\tilde{g}_\delta \leq \int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_{\tilde{g}_\delta} v_\delta|^2 d\tilde{g}_\delta. \quad (61)$$

On the other hand, if we choose w to solve

$$\begin{cases} \Delta_g w = 0 & \text{on } \tilde{M} \setminus \bar{\Omega}_{\delta_0} \\ w = \beta & \text{on } \Sigma \times \{\delta_0\} \\ w(x) \rightarrow 1 & \text{at } \infty, \end{cases} \quad (62)$$

we have that

$$\int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_g w|^2 dg = \lim_{\delta \rightarrow 0} \int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_{\tilde{g}_\delta} w_\delta|^2 d\tilde{g}_\delta, \quad (63)$$

because $\tilde{g}_\delta \rightarrow g$ uniformly on $\tilde{M}$ and $v_\delta \rightarrow \beta$ uniformly on $\Sigma \times \{\delta_0\}$. Hence, it follows from (57), (61) and (63) that

$$\int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_g w|^2 dg \leq \lim_{\delta \rightarrow 0} \int_{\tilde{M} \setminus \bar{\Omega}_{\delta_0}} |\nabla_{\tilde{g}_\delta} v_\delta|^2 d\tilde{g}_\delta = 0, \quad (64)$$

which implies that w must be a constant. Since $\beta < 1$, we get a contradiction. Therefore, $v = 1$ on $\tilde{M} \setminus \bar{\Omega}$.

Next, we let μ, μ_δ denote the $(n-1)$ -dimensional volume measure induced by $g, \tilde{g}_\delta$ on Σ and let e_δ denote the energy $\int_{\tilde{M}} |\nabla_{\tilde{g}_\delta} v_\delta|^2 d\tilde{g}_\delta$.

We fix $0 < \theta < 1$ and $0 < \sigma < \epsilon$. Since $v_\delta \rightarrow 1$ uniformly on compact set away from Σ , we have that

$$v_\delta > \theta \text{ on } \Sigma_\sigma, \text{ for } \delta \ll 1, \quad (65)$$

where Σ_t is the slice $\Sigma \times \{t\}$. We do all the estimates inside the strip $N_\sigma = \Sigma \times [-\sigma, \sigma]$. First, we have that

$$\int_\Sigma \left\{ \int_{-\sigma}^\sigma |\nabla_{\tilde{g}_\delta} v_\delta(x, t)|^2 dt \right\} d\mu_\delta(x) \leq C_1 \int_{N_\sigma} |\nabla_{\tilde{g}_\delta} v_\delta|^2 d\tilde{g}_\delta \leq C_1 e_\delta. \quad (66)$$

Let $l_\delta(x) = \int_{-\sigma}^\sigma |\nabla_{\tilde{g}_\delta} v_\delta(x, t)|^2 dt$, (66) becomes

$$\int_\Sigma l_\delta(x) d\mu_\delta(x) \leq C_1 e_\delta. \quad (67)$$

For any $k > 1, \delta > 0$, we define

$$A_{\delta,k} = \left\{ x \in \Sigma \left| l_\delta(x) \leq k \frac{C_1 e_\delta}{\mu_\delta(\Sigma)} \right. \right\} \quad (68)$$

$$A_{\delta,k}^K = A_{\delta,k} \cap K \quad (69)$$

$$A_{\delta,k,\sigma}^K = A_{\delta,k}^K \times [-\sigma, \sigma]. \quad (70)$$

By (67) we have that

$$\mu_\delta(A_{\delta,k}) \geq \left(1 - \frac{1}{k}\right) \mu_\delta(\Sigma). \quad (71)$$

Since μ_δ is uniformly close to μ , (71) implies that

$$\mu_\delta(A_{\delta,k}^K) \geq \frac{1}{2} \mu_\delta(K) \quad (72)$$

for some fixed large k and any $\delta \ll 1$.

For any $(x, t) \in A_{\delta,k,\sigma}^K$, we have that

$$\begin{aligned} |v_\delta(x, \sigma) - v_\delta(x, t)| &\leq C_2 \int_{-\sigma}^\sigma |\nabla_{\tilde{g}_\delta} v_\delta|(x, t) dt \\ &\leq C_2 (2\sigma)^{\frac{1}{2}} \left\{ \int_{-\sigma}^\sigma |\nabla_{\tilde{g}_\delta} v_\delta|^2(x, t) dt \right\}^{\frac{1}{2}} \\ &= C_2 (2\sigma)^{\frac{1}{2}} l_\delta(x)^{\frac{1}{2}} \\ &\leq C_2 (2\sigma)^{\frac{1}{2}} \left\{ k \frac{C_1 e_\delta}{\mu_\delta(\Sigma)} \right\}^{\frac{1}{2}}. \end{aligned} \quad (73)$$

It follows from (65) that

$$v_\delta(x, t) \geq \theta - C_2(2\sigma)^{\frac{1}{2}} \left\{ k \frac{C_1 e_\delta}{\mu_\delta(\Sigma)} \right\}^{\frac{1}{2}}. \quad (74)$$

On the other hand, for $x \in A_{\delta,k}^K$, we have that

$$\int_{-\delta}^{\delta} \tilde{R}_\delta(x, t) dt \geq u_\delta^{\frac{4}{2-n}}(x) \int_{-\frac{\delta^2}{100}}^{\frac{\delta^2}{100}} \left\{ \eta \left\{ \frac{100}{\delta^2} \phi\left(\frac{100t}{\delta^2}\right) \right\} - C_0 \right\} dt. \quad (75)$$

Therefore, we have the following estimate

$$\begin{aligned} & \liminf_{\delta \rightarrow 0} \int_{A_{\delta,k,\sigma}^K} \tilde{R}_\delta v_\delta^2 d\tilde{g}_\delta \geq \\ & \liminf_{\delta \rightarrow 0} \left\{ \left\{ \theta - C_2 \left\{ (2\sigma) k \frac{C_1 e_\delta}{\mu_\delta(\Sigma)} \right\}^{\frac{1}{2}} \right\}^2 \int_{A_{\delta,k,\sigma}^K} \tilde{R}_\delta d\tilde{g}_\delta \right\} \geq \\ & \theta^2 C_3 \liminf_{\delta \rightarrow 0} \left\{ \int_{A_{\delta,k}^K} \left\{ \int_{-\delta}^{\delta} \tilde{R}_\delta(x, t) dt \right\} d\mu_\delta \right\} \geq \\ & C_3 \theta^2 \eta \liminf_{\delta \rightarrow 0} \mu_\delta(A_{\delta,k}^K) \geq \\ & \frac{1}{2} C_3 \theta^2 \eta \mu(K) > 0 \end{aligned} \quad (76)$$

which is a contradiction to (57). $\square$

We conclude that $\mathcal{G}$ has a strict positive mass in case there exists strict jump of mean curvature across Σ .

5 Zero Mass Case

Let $\mathcal{G} = (g_-, g_+)$ satisfy all the assumptions in Theorem 1 and $m(g_+) = 0$. The following corollary on $R(g_+), R(g_-)$ follows directly from Theorem 1.

Corollary 5.1. *Under the above assumptions, g_- and g_+ both have zero scalar curvature in Ω and $M \setminus \overline{\Omega}$.*

Proof: First, we assume that $R(g_-)$ is not identically zero in Ω . Let u be a positive solution to the equation

$$\begin{cases} \Delta_{g_-} u - c_n R(g_-) u = 0 & \text{on } \Omega \\ u = 1 & \text{on } \Sigma. \end{cases} \quad (77)$$

Consider $\tilde{\mathcal{G}} = (\tilde{g}_-, g_+)$, where $\tilde{g}_- = u^{\frac{4}{n-2}} g_-$. Since u solves the conformal Laplacian of g_- , $\tilde{g}_-$ has zero scalar curvature. By the strong maximum principle, we have $\frac{\partial u}{\partial \nu} > 0$, where ν is the unit outward normal to Σ . A direct computation shows that

$$H(\Sigma, \tilde{g}_-)(x) = H(\Sigma, g_-)(x) + \frac{2}{n-2} \frac{\partial u}{\partial \nu}(x). \quad (78)$$

Hence, $H(\Sigma, \tilde{g}_-) > H(\Sigma, g_-) \geq H(\Sigma, g_+)$. Applying Theorem 1 to $\tilde{\mathcal{G}}$, we see that $m(\tilde{\mathcal{G}}) > 0$, which is a contradiction.

Second, we assume that $R(g_+)$ is not identically zero in $M \setminus \overline{\Omega}$. Let v be a positive solution to

$$\begin{cases} \Delta_{g_+} v - c_n R(g_+) v = 0 & \text{on } M \setminus \overline{\Omega} \\ v = 1 & \text{on } \Sigma \\ v \rightarrow 1 & \text{at } \infty. \end{cases} \quad (79)$$

Consider $\hat{\mathcal{G}} = (g_-, \hat{g}_+)$, where $\hat{g}_+ = v^{\frac{4}{n-2}} g_+$. A similar argument shows that $\hat{g}_+$ is scalar flat in $M \setminus \overline{\Omega}$ and $H(\Sigma, \hat{g}_+) < H(\Sigma, g_+) \leq H(\Sigma, g_-)$. Therefore, Theorem 1 implies that $m(\hat{\mathcal{G}}) > 0$. On the other hand, we have that

$$m(\hat{\mathcal{G}}) = m(\mathcal{G}) + A, \quad (80)$$

where $v = 1 + A|x|^{2-n} + O(|x|^{1-n})$. By the maximum principle, $A \leq 0$. Hence, $m(\mathcal{G}) \geq m(\hat{\mathcal{G}}) > 0$, which is again a contradiction to the assumption that $m(\mathcal{G}) = 0$. $\square$

Corollary 5.1 only reveals information on the scalar curvature, it would be more interesting to know if $m(\mathcal{G}) = 0$ implies that $\mathcal{G}$ is flat away from Σ . Such a type of questions has been studied by H. Bray and F. Finster in [4]. In particular, they obtained the following result concerning the mass and the curvature of a metric which can be approximated by smooth metrics in their sense.

Proposition 5.1. [4] *Suppose $\{g_i\}$ is a sequence of C^3 , complete, asymptotically flat metrics on M^3 with non-negative scalar curvature and the total masses $\{m_i\}$ which converge to a possibly non-smooth limit metric g in the C^0 sense. Let U be the interior of the sets of points where this convergence of metrics is locally C^3 .*

Then if the metrics $\{g_i\}$ have uniformly positive isoperimetric constants and their masses $\{m_i\}$ converges to zero, then g is flat in U .

Now we are in a position to show that, in case $n = 3$, $\mathcal{G}$ is regular cross Σ and $(M, \mathcal{G})$ is isometric to $(\mathbb{R}^3, g_o)$.

Proof of Theorem 2: First, we show that g_- and g_+ are flat in Ω and $M \setminus \bar{\Omega}$. Since g_- and g_+ are $C_{loc}^{3,\alpha}$, it follows from the proof of Proposition 4.1 that $\{\tilde{g}_\delta\}$ converges to g locally in C^3 away from Σ . By Proposition 3.1, we know that $\tilde{g}_\delta$ and g are uniformly close on $\tilde{M}$, hence $\{\tilde{g}_\delta\}$ has uniformly positive isoperimetric constants. By Lemma 4.2, we know that $\lim_{\delta \rightarrow 0} m(\tilde{g}_\delta) = 0$. Therefore, g_- and g_+ are flat by Proposition 5.1.

Second, we show that $A_- = A_+$, where A_- and A_+ are the second fundamental forms of Σ in $(\bar{\Omega}, g_-)$ and $(M \setminus \Omega, g_+)$. Taking trace of the Codazzi equation and using the fact that g_-, g_+ is flat, we have that

$$\begin{cases} \operatorname{div}_{g_\sigma} A_- &= \nabla H(\Sigma, g_-) \\ \operatorname{div}_{g_\sigma} A_+ &= \nabla H(\Sigma, g_+), \end{cases} \quad (81)$$

where g_σ is the induced metric $g_-|_\Sigma = g_+|_\Sigma$. On the other hand, Theorem 1 implies that $H(\Sigma, g_-) \equiv H(\Sigma, g_+)$ on Σ . Hence,

$$\operatorname{div}_{g_\sigma}(A_- - A_+) = 0 \quad \text{and} \quad \operatorname{tr}_{g_\sigma}(A_- - A_+) = 0. \quad (82)$$

We recall the fact that any divergence free and trace free $(0, 2)$ symmetric tensor on (S^2, g_σ) must vanish identically [6], thus we conclude that $A_- = A_+$. Now it follows from the fundamental theorem of surface theory in $\mathbb{R}^3$ that $\mathcal{G}$ is actually C^2 across Σ . The classical PMT [8] then implies that $(M, \mathcal{G})$ is isometric to $\mathbb{R}^3$ with the standard metric. $\square$

Acknowledgments: I would like to thank my Ph.D. advisor Professor Richard Schoen for bringing up this problem and for his superb direction. I also would like to thank Professor Robert Bartnik and Professor Hubert Bray for many stimulating discussions.

References

- [1] Robert Bartnik. New definition of quasilocal mass. *Phys. Rev. Lett.*, 62(20):2346–2348, 1989.
- [2] Robert Bartnik. Energy in general relativity. In *Tsing Hua lectures on geometry & analysis (Hsinchu, 1990–1991)*, pages 5–27. Internat. Press, Cambridge, MA, 1997.
- [3] Hubert Bray. Proof of the riemannian penrose conjecture using the positive mass theorem. *J. Differential Geom.*, 59(2):177–267, 2001.

- [4] Hubert Bray and Felix Finster. Curvature estimates and the positive mass theorem. *Comm. Anal. Geom.*, 10(2):291–306, 2002.
- [5] Gilbarg David and Trudinger Neil S. *Elliptic Partial Differential Equations of Second Order*. Berlin: Springer-Verlag, 1983.
- [6] Heinz Hopf. *Differential geometry in the large*, volume 1000 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, second edition, 1989. Notes taken by Peter Lax and John W. Gray, With a preface by S. S. Chern, With a preface by K. Voss.
- [7] Richard Schoen. Variational theory for the total scalar curvature functional for riemannian metrics and related topics. In *Topics in the Calculus of Variations. Lecture Notes in Math. 1365*, pages 120–154. Berlin: Springer-Verlag, 1987.
- [8] Richard Schoen and Shing Tung Yau. On the proof of the positive mass conjecture in general relativity. *Comm. Math. Phys.*, 65(1):45–76, 1979.